



ARSET Applied Remote Sensing Training http://arset.gsfc.nasa.gov

@NASAARSET

Satellite Based Fire Products:
Methods, Data Access, and Applications

Melanie Follette-Cook, Pawan Gupta, & Land Team


Satellite Remote Sensing of Air Quality


September 19-21, 2017 University of California, Riverside

www.nasa.gov

MODIS


- Spatial Resolution
 - -250 m, 500 m, 1 km
- Temporal Resolution
 - Daily, 8 day, 16 day, monthly, quarterly, yearly
 - -2000-present
- Data Format
 - Hierarchal data format EarthObserving System Format (HDF–EO8)
- Spectral Coverage
 - –36 bands (major bands include red, blue, IR, NIR, MIR)
 - Bands 1-2: 250 m
 - Bands 3-7: 500 m
 - Bands 8-36: 1000 m





MODIS Fire Products

- Near Real-Time (NRT) thermal anomalies and fire locations
- Provides snapshots of active burning fires and burned areas
- The Active Fire product delivers actively burning locations on a daily basis at 1 km resolution (additional 8 day and monthly products)



Global Fire Map (September 17 – 26, 2016)

Colors range from red, where the fire count is low, to yellow where the number of fires is large

MODIS Thermal Anomalies Algorithm

- MODIS Fire Detection:
 - -1 km pixel flagged as containing one or more fires
 - –can also detect volcanic signatures
- Significant increase in absolute radiance at 4 μm (band 22) and 11 μm (band 31)
 - -cloud masks applied
 - VIIRS active fire detection algorithm is similar



VIIRS fire detections, NASA Worldview

MODIS Fire Detection Algorithm

http://modis-fire.umd.edu/files/atbd_mod14.pdf


Table 2: MODIS channels used for active-fire detection and characterization.

| | Central | |
|---------|-----------------|---|
| Channel | wavelength (µm) | Purpose |
| 1 | 0.65 | Sun glint and coastal false alarm rejection; cloud masking. |
| 2 | 0.86 | Bright surface, sun glint, and coastal false alarm rejection; |
| | | cloud masking. |
| 7 | 2.1 | Sun glint and coastal false alarm rejection. |
| 21 | 3.96 | High-range channel for fire detection and characterization. |
| 22 | 3.96 | Low-range channel for fire detection and characterization. |
| 31 | 11.0 | Fire detection, cloud masking. |
| 32 | 12.0 | Cloud masking. |

- Potential fire pixel identified
 - -BT4 > 310 K
 - -BT4 BT11 > 10 K
 - -0.86 reflectance < 0.3
- Otherwise flagged as non-fire pixel

MODIS Thermal Anomalies Algorithm


- Limitations
 - -False positives: small forest clearings (bare soil)
 - Large fire omissions due thick smoke
- Collection 6 (most recent) improves upon these errors
 - –Global commission error of 1.2%



MODIS fire detections, NASA Worldview

MODIS Land Products: Burned Area (MCD64A1)

- The combined Terra & Aqua MODIS Burned Area Product is a monthly gridded 500m product
- MODIS detects the approximate date of burning at 500m resolution
- Maps include the spatial extent of recent fires
- For more information: http://modis-fire.umd.edu



This image shows the extent of the Long Draw fire that occurred in southeastern Oregon The colors represent the approximate day of the burning from July 8 (start of fire) to July 12, 2012 (end of fire)

Where to Obtain MODIS Products



Land Process Distributed Active Archive (LPDAAC) http://lpdaac.usgs.gov/



ECHO Reverb: http://reverb.echo.nasa.gov



Worldview:


http://worldview.earthdata.nasa.gov



Fire Information for Resource Management System: http://earthdata.nasa.gov/earth- observation-data/near-real-time/firms


Fire Information for Resource Management System (FIRMS)

- Near real-time (NRT) active fire data within 3 hrs of satellite overpass
- Global MODIS and VIIRS hotspots, fire locations, and burned area images
- Historical data available
- Available in:
 - -Email alerts
 - -GIS-friendly file format
 - Visualization in Web Fire Mapper or Worldview



VIIRS Active Fire Product


- Released October 22, 2012
- Spatial resolution:
 - -750 m (M-band)
 - -375 m (I-band)
- Data still preliminary and continually undergo evaluation & calibration
- Data available as:
 - -ASCII
 - -KMZ
 - -TIFF
- Exercise on this tool in upcoming session



Northern California Fires, 2015


VIIRS Active Fire Map (CONUS)

http://viirsfire.geog.umd.edu/map/viirsMap.php



VIIRS Active Fire Map (CONUS)

http://viirsfire.geog.umd.edu/map/map_v2.php



US Forest Service - Tools

https://fsapps.nwcg.gov/afm/




Where to Obtain VIIRS Land Products



Worldview:

http://worldview.earthdata.nasa.gov



VIIRS Active Fire:

http://viirsfire.geog.umd.edu/pages/about.php



NOAA Comprehensive Large Array-Data Stewardship System (CLASS):

http://www.class.ngdc.noaa.gov/saa/products/welcome



Level-1 and Atmosphere Archive & Distribution System Website: http://ladsweb.nascom.nasa.gov

Questions & Discussion Prompts

- Changes in what retrieved quantity are used to detect fires?
- What is a source of uncertainty for fire detection?

